

Multiple Choice (5 points each, Put answers in CAPS in the left margin.)

$$R = 8.314 \text{ J/mol}\cdot\text{K} = 0.0821 \text{ L}\cdot\text{atm/mol}\cdot\text{K}$$

- The additional energy added to a solid to cause it to liquefy is called its
 - Heat of formation
 - Lattice energy
 - Heat of fusion
 - Heat of vaporization
 - Heat of liquefaction
- Which of the following has the strongest solute-to-solvent interaction? (Problem 13.4)
 - NH_3 in H_2O
 - CH_4 in C_6H_6
 - KCl in H_2O
 - CH_2Cl_2 in C_6H_6
 - CH_2Cl_2 in CH_3OH
- For the compounds CH_4 , CH_3Cl , and CH_3OH , what is the correct trend with respect to their boiling points? (Homework 10.10)
 - $\text{CH}_4 < \text{CH}_3\text{Cl} < \text{CH}_3\text{OH}$
 - $\text{CH}_4 < \text{CH}_3\text{OH} < \text{CH}_3\text{Cl}$
 - $\text{CH}_3\text{Cl} < \text{CH}_4 < \text{CH}_3\text{OH}$
 - $\text{CH}_3\text{Cl} < \text{CH}_3\text{OH} < \text{CH}_4$
 - $\text{CH}_3\text{OH} < \text{CH}_4 < \text{CH}_3\text{Cl}$
 - $\text{CH}_3\text{OH} < \text{CH}_3\text{Cl} < \text{CH}_4$
- Which of the following pairs is unlikely to be miscible?
 - CH_3OH and H_2O
 - CH_3OH and CH_3Br
 - CH_3Br and H_2O
 - CH_3OH and $\text{CH}_3\text{CH}_2\text{OH}$
 - H_2O and $\text{CH}_3(\text{CH}_2)_4\text{CH}_3$
- Which of the following would have the highest boiling point? (Homework 11.24)
 - 0.12 *m* KI
 - 0.35 *m* $\text{C}_2\text{H}_5\text{OH}$
 - 0.30 *m* NH_3
 - 0.19 *m* CaF_2
 - pure H_2O
- Which of the following is false about colloids?
 - They are generally opaque or translucent.
 - Brownian motion accounts for the existence and appearance of colloids.
 - They are usually composed of very small solid particles suspended in a liquid.
 - Filtration is an easy way to separate a colloid.
 - Milk, mayonnaise, and shaving cream are common colloids.
- What is the reaction order for the following elementary reaction: $\text{A} + \text{B} \rightarrow \text{C} + \text{D}$
 - 1
 - 2
 - 3
 - 4
 - cannot determine from given information

8. The reaction $2 \text{NO}_2(\text{g}) \longrightarrow 2 \text{NO}(\text{g}) + \text{O}_2(\text{g})$ is suspected to be second order in NO_2 . Which of the following kinetic plots would be the best to do to prove the reaction is second order?

A) $[\text{NO}_2]$ vs t

C) $[\text{NO}_2]^2$ vs t

E) $\ln[\text{NO}_2]^{-1}$ vs t

B) $\ln[\text{NO}_2]$ vs t

D) $[\text{NO}_2]^{-1}$ vs t

Discussion Questions (You must show your work to receive credit):

1. At 1 atm, how much energy is required to heat 57.0 g $\text{H}_2\text{O}_{(\text{s})}$ at -20.0°C to $\text{H}_2\text{O}_{(\text{l})}$ at 59.0°C ? Specific heat $\text{H}_2\text{O}_{(\text{s})} = 2.087 \text{ J}/(\text{g}\cdot^\circ\text{C})$ and specific heat $\text{H}_2\text{O}_{(\text{l})} = 4.184 \text{ J}/(\text{g}\cdot^\circ\text{C})$, $\Delta H_{\text{fus}}^\circ = 6.010 \text{ kJ/mol}$ (8 points)

2. Define viscosity and explain how it changes with changing intermolecular forces. (8 points)

3. At 298 K, the Henry's law constant for oxygen is 0.00130 M/atm. Air is 21.0% oxygen. At 298 K, what is the solubility of oxygen in water exposed to pure oxygen? To air at 1.00 atm? (10 points)

4. A solution of H_2SO_4 (aq) with a molal concentration of 2.24 *m* has a density of 1.135 g/mL. What is the molar concentration of this solution? (5 points)
5. One mole of KCl is added to a liter of pure water and one mole of MgF_2 is added to a different liter of pure water. Which has the higher boiling point? Explain your answer (5 points)
5. For the reaction $2 \text{N}_2\text{O}_5(\text{g}) \longrightarrow 4 \text{NO}_2(\text{g}) + \text{O}_2(\text{g})$, the activation energy and overall ΔE are 100 kJ/mol and -23 kJ/mol respectively. (a) Sketch the energy profile for this reaction. (b) What is the activation energy for the reverse reaction? (9 points)

6. For the reaction $\text{C}_5\text{H}_{10} + \text{O}_3 \longrightarrow \text{C}_5\text{H}_{10}\text{O}_3$ the following data were collected:

$[\text{C}_5\text{H}_{10}]$ (M)	$[\text{O}_3]$ (M)	rate (Ms^{-1})
0.010	0.0028	2.2
0.0050	0.0028	1.1
0.010	0.010	4.4

- a) What is the rate law for this reaction?
- b) What is the value of the rate constant for this reaction? (15 points)
- c) Could this be an elementary reaction? Explain.